

UBE4B Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID AAD02233.1

Catalog No. A301-124A


GeneID 10277

Lot No. A301-124A-1



APPLICATIONS	IP
SPECIES REACTIVITY	Human
AMOUNT	100 µl
CONCENTRATION	1000 µg/ml
STORAGE/SHELF LIFE	2 – 8° C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antibody was affinity purified using an epitope specific to UBE4B immobilized on solid support. The epitope recognized by A301-124A maps to a region between residue 1252 and 1302 of human ubiquitination factor E4B using the numbering given in entry AAD02233.1 (GeneID 10277). Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.
APPLICATIONS	Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot Not recommended. Use rabbit anti-UBE4B antibody A301-123A. Immunoprecipitation 2 – 5 µg/mg lysate
APPLICATION NOTES	Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).
ADDITIONAL INFO	https://www.bethyl.com/product/A301-124A Use the link above to view SDS, a current list of citations, and other product specific information. IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019



Detection of human UBE4B by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1 mg for IP, 20% of IP loaded) from HeLa cells. *Antibodies:* Affinity purified rabbit anti-UBE4B antibody A301-124A used for IP at 3 µg/mg lysate. UBE4B was also immunoprecipitated by rabbit anti-UBE4B antibody A301-123A, which recognizes an upstream epitope. For blotting immunoprecipitated UBE4B, A301-123A was used at 0.1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 3 minutes.