APC4 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_037499.2

Catalog No. A301-175A GeneID 29945

Lot No. A301-175A-1

APPLICATIONS IP

SPECIES REACTIVITY Human

PRESUMED REACTIVITY Based on 100% sequence identity, this antibody is predicted to react with Mouse and Orangutan

AMOUNT 100 μl

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to APC4 immobilized on solid support.

The epitope recognized by A301-175A maps to a region between residue 425 and 475 of human anaphase-promoting complex subunit 4 using the numbering given in entry NP_037499.2

(GeneID 29945).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm

of 1.4 equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot Not recommended. Use rabbit anti-APC4 antibody A301-176A.

Immunoprecipitation $2 - 5 \mu g/mg$ lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100-020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

ADDITIONAL INFO https://www.bethyl.com/product/A301-175A


Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019



APC4 Antibody A301-175A



Detection of human APC4 by western blot of immunoprecipitates. Samples: Whole cell lysate (1 mg for IP, 20% of IP loaded) from HeLa cells. Antibodies: Affinity purified rabbit anti-APC4 antibody A301-175A used for IP at 3 μg/mg lysate. APC4 was also immunoprecipitated by rabbit anti-APC4 antibody A301-176A, which recognizes a downstream epitope. For blotting immunoprecipitated APC4, A301-176A was used at 1.0 μg/ml. Detection: Chemiluminescence with an exposure time of 1 second.