TRIM47/GOA Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_258411.2

Catalog No. A301-194A GenelD 91107

Lot No. A301-194A-1

APPLICATIONS IP

SPECIES REACTIVITY Human

AMOUNT 100 μl

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION Antibody was affinity purified using an epitope specific to TRIM47/GOA immobilized on solid

PROCEDURES support.

The epitope recognized by A301-194A maps to a region between residue 400 and 450 of human tripartite motif-containing 47 (gene overexpressed in astrocytoma) using the numbering given in

entry NP_258411.2 (GeneID 91107).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm

of 1.4 equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot Not recommended. Use rabbit anti-TRIM47/GOA antibody A301-

193A.

Immunoprecipitation $2 - 5 \mu g/mg$ lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100-020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

ADDITIONAL INFO https://www.bethyl.com/product/A301-194A


Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019





Detection of human TRIM47/GOA by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1 mg for IP, 20% of IP loaded) from HeLa cells. *Antibodies:* Affinity purified rabbit anti–TRIM47/GOA antibody A301–194A used for IP at 3 μ g/mg lysate. TRIM47/GOA was also immunoprecipitated by rabbit anti–TRIM47/GOA antibody A301–193A, which recognizes an upstream epitope. For blotting immunoprecipitated TRIM47/GOA, A301–193A was used at 1 μ g/ml. *Detection:* Chemiluminescence with an exposure time of 3 seconds.