## AktS1/PRAS40 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP\_115751.1

Catalog No. A301-201A GenelD 84335

Lot No. A301-201A-1

APPLICATIONS WB, IP
SPECIES REACTIVITY Human

PRESUMED REACTIVITY Based on 100% sequence identity, this antibody is predicted to react with Mouse

**AMOUNT** 100 μl

CONCENTRATION 200 μg/ml

**STORAGE/SHELF LIFE** 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

**BUFFER** Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

**PRODUCTION** Antibody was affinity purified using an epitope specific to AktS1/PRAS40 immobilized on solid

**PROCEDURES** support.

The epitope recognized by A301-201A maps to a region between residue 195 and 245 of human

AKT1 substrate 1 (proline-rich Akt substrate, 40 kDa) using the numbering given in entry

NP 115751.1 (GenelD 84335).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm

of 1.4 equals 1.0 mg of IgG.

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 - 1:10,000

Immunoprecipitation  $2 - 5 \mu g/mg$  lysate

**APPLICATION NOTES** Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100-020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-20% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-20% SDS-PAGE.


ADDITIONAL INFO https://www.bethyl.com/product/A301-201A

Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol\_IP\_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019





Detection of human AktS1/PRAS40 by western blot and immunoprecipitation. Samples: Whole cell lysate from HeLa (5, 15 and 50 μg for WB; 1 mg for IP, 20% of IP loaded) and HEK293T (T; 50 μg) cells. Antibodies: Affinity purified rabbit anti-AktS1/PRAS40 antibody A301-201A used for WB at 0.04 μg/ml (A) and 1 μg/ml (B) and used for IP at 3 μg/mg lysate. AktS1/PRAS40 was also immunoprecipitated by rabbit anti-AktS1/PRAS40 antibody A301-202A, which recognizes a downstream epitope. Detection: Chemiluminescence with exposure times of 30 seconds (A) and 10 seconds (B).