

JHDM1A Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID NP_036440.1

Catalog No. A301-476A

GeneID 22992

Lot No. A301-476A-2



APPLICATIONS	WB, IP
SPECIES REACTIVITY	Human, Mouse
AMOUNT	100 µl
CONCENTRATION	200 µg/ml
STORAGE/SHELF LIFE	2 – 8° C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antibody was affinity purified using an epitope specific to JHDM1A immobilized on solid support.

The epitope recognized by A301-476A maps to a region between residue 1112 and 1162 of human JmjC domain-containing histone demethylation protein 1A using the numbering given in entry NP_036440.1 (GeneID 22992).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 – 1:10,000

Immunoprecipitation 2 – 10 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).


Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.

ADDITIONAL INFO <https://www.bethyl.com/product/A301-476A>


Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019



Detection of human and mouse JHDM1A by western blot.
Samples: Whole cell lysate (50 µg) from HeLa, HEK293T, and mouse NIH 3T3 cells prepared using NETN lysis buffer.
Antibodies: Affinity purified rabbit anti-JHDM1A antibody A301-476A (lot A301-476A-2) used for WB at 0.2 µg/ml.
Detection: Chemiluminescence with an exposure time of 3 minutes.



Detection of human JHDM1A by western blot of immunoprecipitates.
Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from HeLa cells prepared using NETN lysis buffer.
Antibodies: Affinity purified rabbit anti-JHDM1A antibody A301-476A (lot A301-476A-2) used for IP at 6 µg per reaction. JHDM1A was also immunoprecipitated by a previous lot of this antibody (lot A301-476A-1). For blotting immunoprecipitated JHDM1A, A301-476A was used at 1 µg/ml.
Detection: Chemiluminescence with an exposure time of 3 minutes.