

USP11 Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID EAW59289.1

Catalog No. A301-613A


GeneID 8237

Lot No. A301-613A-2




APPLICATIONS	WB, IP
SPECIES REACTIVITY	Human
AMOUNT	100 µl
CONCENTRATION	200 µg/ml
STORAGE/SHELF LIFE	2 – 8° C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	<p>Antibody was affinity purified using an epitope specific to USP11 immobilized on solid support.</p> <p>The epitope recognized by A301-613A maps to a region between residue 870 and 920 of human ubiquitin specific peptidase 11 using the numbering given in entry EAW59289.1 (GeneID 8237).</p> <p>Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.</p>
APPLICATIONS	<p>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.</p> <p>Western Blot 1:2,000 – 1:10,000</p> <p>Immunoprecipitation 2 – 10 µg/mg lysate</p>
APPLICATION NOTES	<p>Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).</p> <p>Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.</p>
ADDITIONAL INFO	<p>https://www.bethyl.com/product/A301-613A</p> <p>Use the link above to view SDS, a current list of citations, and other product specific information.</p> <p>IP-western blot protocol https://www.bethyl.com/content/protocol_IP_WB</p>

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Eric McIntush, PhD | Chief Scientific Officer Date: December 13, 2019



Detection of human USP11 by western blot. *Samples:* Whole cell lysate (50 µg) from Jurkat, Hep-G2, U2OS, HEK293T, and MCF-7 cells prepared using NETN lysis buffer. *Antibody:* Affinity purified rabbit anti-USP11 antibody A301-613A (lot A301-613A-2) used for WB at 0.04 µg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.



Detection of human USP11 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from Jurkat cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-USP11 antibody A301-613A (lot A301-613A-2) used for IP at 6 µg per reaction. USP11 was also immunoprecipitated by a previous lot of this antibody (lot A301-613A-1). For blotting immunoprecipitated USP11, A301-613A was used at 0.04 µg/ml. *Detection:* Chemiluminescence with an exposure time of 1 second.