


Detection of human TReP-132/TRERF1 by western blot.


Samples: Whole cell lysate (50 µg) from HeLa, HEK293T, and Jurkat cells prepared using NETN lysis buffer.

Antibody: Affinity purified rabbit anti-TReP-132/TRERF1 antibody A301-631A (lot A301-631A-2) used for WB at 0.1 µg/ml. **Detection:** Chemiluminescence with an exposure time of 3 minutes.



Detection of human TReP-132/TRERF1 by western blot of immunoprecipitates.

Samples: Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from HEK293T cells prepared using NETN lysis buffer. **Antibodies:** Affinity purified rabbit anti-TReP-132/TRERF1 antibody A301-631A (lot A301-631A-2) used for IP at 3 µg per reaction. TReP-132/TRERF1 was also immunoprecipitated by rabbit anti-TReP-132/TRERF1 antibody A301-630A. For blotting immunoprecipitated TReP-132/TRERF1, A301-631A was used at 1 µg/ml. **Detection:** Chemiluminescence with an exposure time of 3 minutes.



Detection of human TReP-132/TRERF1 by immunohistochemistry. **Sample:** FFPE section of human ovarian carcinoma. **Antibody:** Affinity purified rabbit anti-TReP-132/TRERF1 (Cat. No. A301-631A Lot2) used at a dilution of 1:5,000 (0.2 µg/ml). **Detection:** DAB