## **BCoR Antibody**

Rabbit Polyclonal

Antigen Affinity Purified Protein ID EAW59426.1

Catalog No. A301-672A GeneID 54880

Lot No. A301-672A-2

APPLICATIONS IP

SPECIES REACTIVITY Human

AMOUNT 100 μl

CONCENTRATION 1000 μg/ml

**STORAGE/SHELF LIFE** 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

**BUFFER** Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to BCoR immobilized on solid support.

The epitope recognized by A301-672A maps to a region between residue 25 and 75 of human

BCL6 co-repressor using the numbering given in entry EAW59426.1 (GenelD 54880).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm

of 1.4 equals 1.0 mg of IgG.

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot Not recommended. Use rabbit anti-BCoR antibody A301-673A.

Immunoprecipitation  $2 - 10 \mu g/mg$  lysate

**APPLICATION NOTES** Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100–020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

ADDITIONAL INFO https://www.bethyl.com/product/A301-672A

Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol\_IP\_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019



BCoR Antibody A301-672A



Detection of human BCoR by western blot of immunoprecipitates. Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from HeLa cells prepared using NETN lysis buffer. Antibodies: Affinity purified rabbit anti-BCoR antibody A301-672A (lot A301-672A-2) used for IP at 6 μg per reaction. BCoR was also immunoprecipitated by rabbit anti-BCoR antibody A301-672A-1. For blotting immunoprecipitated BCoR, A301-673A was used at 1 μg/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.