## SAS-6 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP\_919268.1

Catalog No. A301-802A GenelD 163786

Lot No. A301-802A-1

APPLICATIONS WB, IP

SPECIES REACTIVITY Human

AMOUNT 100 μI

 $\textbf{CONCENTRATION} \qquad \quad 1000 \; \mu\text{g/ml}$ 

**STORAGE/SHELF LIFE** 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to SAS-6 immobilized on solid support.

The epitope recognized by A301-802A maps to a region between residue 607 and 657 of human

spindle assembly 6 using the numbering given in entry NP\_919268.1 (GeneID 163786).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm

of 1.4 equals 1.0 mg of IgG.

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 - 1:10,000 Immunoprecipitation 2 - 5 µg/mg lysate

**APPLICATION NOTES** Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100–020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.

ADDITIONAL INFO https://www.bethvl.com/product/A301-802A


Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol IP WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019



SAS-6 Antibody A301-802A



Detection of human SAS-6 by western blot and immunoprecipitation. Samples: Whole cell lysate from HeLa (5, 15 and 50  $\mu$ g for WB; 1 mg for IP, 20% of IP loaded) and HEK293T (T; 50  $\mu$ g) cells. Antibody: Affinity purified rabbit anti–SAS-6 antibody A301–802A used for WB at 0.1  $\mu$ g/ml (A) and 1  $\mu$ g/ml (B) and used for IP at 3  $\mu$ g/mg lysate. SAS-6 was also immunoprecipitated by rabbit anti–SAS-6 antibody A301–801A, which recognizes an upstream epitope. Detection: Chemiluminescence with exposure times of 3 minutes (A) and 30 seconds (B).