CCDC6 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID AAC60637.1

Catalog No. A302-097A GeneID 8030

Lot No. A302-097A-1

APPLICATIONS IP

SPECIES REACTIVITY Human

AMOUNT 100 μl

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to CCDC6 immobilized on solid support.

The epitope recognized by A302–097A maps to a region between residue 375 and 425 of human coiled-coil domain containing 6 using the numbering given in entry AAC60637.1 (GeneID 8030).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm

of 1.4 equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot Not recommended. Use rabbit anti-CCDC6 antibody A302-096A.

Immunoprecipitation 2 – 5 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100–020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-20% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

ADDITIONAL INFO https://www.bethyl.com/product/A302-097A

Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019



CCDC6 Antibody



Detection of human CCDC6 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1 mg for IP, 20% of IP loaded) from HeLa cells. *Antibodies:* Affinity purified rabbit anti-CCDC6 antibody A302-097A used for IP at 3 μg/mg lysate. CCDC6 was also immunoprecipitated by rabbit anti-CCDC6 antibody A302-096A, which recognizes an upstream epitope. For blotting immunoprecipitated CCDC6, A302-096A was used at 1 μg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.