GAPex5 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID BAG10470.1

Catalog No. A302-116A GeneID 26130

Lot No. A302-116A-1

APPLICATIONS WB, IP
SPECIES REACTIVITY Human

PRESUMED REACTIVITY Based on 100% sequence identity, this antibody is predicted to react with Mouse and Bovine

AMOUNT 100 μl

CONCENTRATION 200 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to GAPex5 immobilized on solid support.

The epitope recognized by A302-116A maps to a region between residue 700 and 750 of human GTPase activating protein and VPS9 domains 1 using the numbering given in entry BAG10470.1

(GeneID 26130).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm

of 1.4 equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 - 1:10,000

Immunoprecipitation 2 – 5 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100-020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.

ADDITIONAL INFO https://www.bethyl.com/product/A302-116A


Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019





Detection of human GAPex5 by western blot and immunoprecipitation. *Samples:* Whole cell lysate (5, 15 and 50 μg for WB; 1 mg for IP, 20% of IP loaded) from HeLa cells. *Antibodies:* Affinity purified rabbit anti-GAPex5 antibody A302–116A used for WB at 0.04 μg/ml (A) and 0.1 μg/ml (B) and used for IP at 3 μg/mg lysate. GAPex5 was also immunoprecipitated by rabbit anti-GAPex5 antibody A302–115A, which recognizes an upstream epitope. *Detection:* Chemiluminescence with exposure times of 10 seconds (A and B).