## PIK3R2 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP\_005018.1

Catalog No. A302-594A GeneID 5296

Lot No. A302-594A-1

APPLICATIONS IP

**SPECIES REACTIVITY** Human 100 μl

CONCENTRATION 1000 µg/ml

**STORAGE/SHELF LIFE** 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to PIK3R2 immobilized on solid support.

The epitope recognized by A302-594A maps to a region between residue 125 and 175 of human

phosphoinositide-3-kinase, regulatory subunit 2 using the numbering given in entry

NP\_005018.1 (GeneID 5296).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot Not recommended. Use rabbit anti-PIK3R2 antibody A302-593A.

Immunoprecipitation 5 – 15 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

ADDITIONAL INFO https://www.bethyl.com/product/A302-594A


Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol\_IP\_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019





Detection of human PIK3R2 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1 mg for IP, 20% of IP loaded) from HeLa cells. *Antibodies:* Affinity purified rabbit anti-PIK3R2 antibody A302-594A used for IP at 10 μg/mg lysate. PIK3R2 was also immunoprecipitated by rabbit anti-PIK3R2 antibody A302-593A, which recognizes an upstream epitope. For blotting immunoprecipitated PIK3R2, A302-593A was used at 1 μg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.