

# TNIK Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID NP\_055843.1

Catalog No. A302-696A


GeneID 23043

Lot No. A302-696A-2



<b>APPLICATIONS</b>	IP
<b>SPECIES REACTIVITY</b>	Human
<b>AMOUNT</b>	100 $\mu$ l
<b>CONCENTRATION</b>	1000 $\mu$ g/ml
<b>STORAGE/SHELF LIFE</b>	2 - 8° C / 1 year from date of receipt
<b>PHYSICAL STATE</b>	Liquid
<b>BUFFER</b>	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
<b>ISOTYPE</b>	IgG
<b>ORIGIN</b>	USA
<b>PRODUCTION PROCEDURES</b>	<p>Antibody was affinity purified using an epitope specific to TNIK immobilized on solid support.</p> <p>The epitope recognized by A302-696A maps to a region between residue 750 and 800 of human TRAF2 and NCK interacting kinase using the numbering given in entry NP_055843.1 (GeneID 23043).</p> <p>Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.</p>
<b>APPLICATIONS</b>	<p>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.</p> <p>Western Blot                      Not recommended. Use rabbit anti-TNIK antibody A302-695A.</p> <p>Immunoprecipitation              2 - 10 <math>\mu</math>g/mg lysate</p>
<b>APPLICATION NOTES</b>	Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).
<b>ADDITIONAL INFO</b>	<p><a href="https://www.bethyl.com/product/A302-696A">https://www.bethyl.com/product/A302-696A</a></p> <p>Use the link above to view SDS, a current list of citations, and other product specific information. IP-western blot protocol: <a href="https://www.bethyl.com/content/protocol_IP_WB">https://www.bethyl.com/content/protocol_IP_WB</a></p>

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.  
Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019



**Detection of human TNIK by western blot of immunoprecipitates.** *Samples:* Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from HeLa cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-TNIK antibody A302-696A (lot A302-696A-2) used for IP at 6 µg per reaction. TNIK was also immunoprecipitated by a previous lot of this antibody (lot A302-696A-1) and rabbit anti-TNIK antibody A302-695A. For blotting immunoprecipitated TNIK, A302-696A was used at 1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 3 minutes.