USP8 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_005145.3

Catalog No. A302-930A GenelD 9101

Lot No. A302-930A-1

APPLICATIONS IP

SPECIES REACTIVITY Human 100 μl

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to USP8 immobilized on solid support.

The epitope recognized by A302-930A maps to a region between residue 1068 and 1118 of human Ubiquitin Specific Peptidase 8 using the numbering given in entry NP_005145.3 (GeneID

9101).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot Not recommended

Immunoprecipitation 2 – 5 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

ADDITIONAL INFO https://www.bethyl.com/product/A302-930A

Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019



USP8 Antibody A302-930A



Detection of human USP8 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1 mg for IP, 20% of IP loaded) from HeLa cells. *Antibodies:* Affinity purified rabbit anti–USP8 antibody A302–930A used for IP at 3 μg/mg lysate. USP8 was also immunoprecipitated by rabbit anti–USP8 antibody A302–929A, which recognizes an upstream epitope. For blotting immunoprecipitated USP8, A302–929A was used at 0.4 μg/ml,. *Detection:* Chemiluminescence with an exposure time of 10 seconds.