

HEXIM1 Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID NP_006451.1

Catalog No. A303-113A



GeneID 10614

Lot No. A303-113A-1




| | |
|------------------------------|---|
| APPLICATIONS | WB, IP, IHC |
| SPECIES REACTIVITY | Human, Mouse |
| AMOUNT | 100 µl |
| CONCENTRATION | 200 µg/ml |
| STORAGE/SHELF LIFE | 2 - 8° C / 1 year from date of receipt |
| PHYSICAL STATE | Liquid |
| BUFFER | Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide |
| ISOTYPE | IgG |
| ORIGIN | USA |
| PRODUCTION PROCEDURES | <p>Antibody was affinity purified using an epitope specific to HEXIM1 immobilized on solid support.</p> <p>The epitope recognized by A303-113A maps to a region between residue 309 and 359 of human Hexamethylene Bis-Acetamide Inducible 1 using the numbering given in entry NP_006451.1 (GeneID 10614).</p> <p>Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.</p> |
| APPLICATIONS | <p>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.</p> <p>Western Blot 1:2,000 - 1:10,000</p> <p>Immunoprecipitation 2 - 10 µg/mg lysate</p> <p>Immunohistochemistry 1:200 - 1:1,000. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections.</p> |
| APPLICATION NOTES | <p>Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-20% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).</p> <p>Western blot of lysates performed using standard western blot reagents and 4-20% SDS-PAGE.</p> |
| IHC HUMAN CONTROLS | Breast Carcinoma, Ovarian Carcinoma, Prostate Carcinoma, Stomach Adenocarcinoma |
| IHC MOUSE CONTROLS | Teratoma |
| ADDITIONAL INFO | <p>https://www.bethyl.com/product/A303-113A</p> <p>Use the link above to view SDS, a current list of citations, and other product specific information.</p> <p>IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB</p> |

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019



Detection of human HEXIM1 by western blot and immunoprecipitation. *Samples:* Whole cell lysate (5, 15 and 50 µg for WB; 1 mg for IP, 20% of IP loaded) from HeLa cells. *Antibodies:* Affinity purified rabbit anti-HEXIM1 antibody A303-113A used for WB at 0.04 µg/ml (A) and 1 µg/ml (B) and used for IP at 6 µg/mg lysate. HEXIM1 was also immunoprecipitated by rabbit anti-HEXIM1 antibodies A303-111A and A303-112A, which recognize upstream epitopes. *Detection:* Chemiluminescence with exposure times of 30 seconds (A) and 3 seconds (B).

Detection of human HEXIM1 by immunohistochemistry. *Samples:* FFPE serial sections of human lung cancer. *Antibody:* Affinity purified rabbit anti-HEXIM1 (Cat. No. A303-111A Lot1, left image, A303-112A Lot1, middle image and Cat. No. A303-113A Lot1, right image) used at a dilution of 1:200 (1 µg/ml) (A303-111A-1) or 1:1,000 (1 µg/ml) (A303-112A-1 and A303-113A-1). *Detection:* DAB. *Counterstain:* hematoxylin (blue).



Detection of mouse HEXIM1 by immunohistochemistry. *Sample:* FFPE section of mouse teratoma. *Antibody:* Affinity purified rabbit anti-HEXIM1 (Cat. No. A303-113A Lot1) used at a dilution of 1:200 (1 µg/ml). *Detection:* DAB