HOXC10 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_059105.2

Catalog No. A303-177A GeneID 3226

Lot No. A303-177A-1

APPLICATIONS IP

SPECIES REACTIVITY Human 100 μl

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to HOXC10 immobilized on solid support.

The epitope recognized by A303-177A maps to a region between residue 50 and 100 of human

Homeobox C10 using the numbering given in entry NP_059105.2 (GeneID 3226).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot Not recommended

Immunoprecipitation 2 – 10 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100–020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-20% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

ADDITIONAL INFO https://www.bethyl.com/product/A303-177A


Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019





Detection of human HOXC10 by western blot of immunoprecipitates. Samples: Whole cell lysate from HeLa (1 mg for IP, 20% of IP loaded) cells. Antibodies: Affinity purified rabbit anti-HOXC10 antibody A303-177A used for IP at 6 μ g/mg lysate. HOXC10 was also immunoprecipitated by rabbit anti-HOXC10 antibody A303-178A, which recognizes a downstream epitope. For blotting immunoprecipitated HOXC10, A303-178A was used at 1 μ g/ml. Detection: Chemiluminescence with an exposure time of 10 seconds.