ACBD5 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID CAI16912.1

Catalog No. A303-296A GeneID 91452

Lot No. A303-296A-1

APPLICATIONS IP

SPECIES REACTIVITY Human

AMOUNT 100 μl

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to ACBD5 immobilized on solid support.

The epitope recognized by A303-296A maps to a region between residue 325 and 375 of human Acyl-CoA Binding Domain Containing 5 using the numbering given in entry CAI16912.1 (GenelD

91452).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot Not recommended

Immunoprecipitation 2 – 10 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-20% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

ADDITIONAL INFO https://www.bethyl.com/product/A303-296A

Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019



ACBD5 Antibody



Detection of human ACBD5 by western blot of immunoprecipitates. Samples: Whole cell lysate (1 mg for IP, 20% of IP loaded) from HeLa cells. Antibodies: Affinity purified rabbit anti-ACBD5 antibody A303-296A used for IP at 6 μ g/mg lysate. ACBD5 was also immunoprecipitated by rabbit anti-ACBD5 antibody A303-297A, which recognizes a downstream epitope. For blotting immunoprecipitated ACBD5, A303-297A was used at 1 μ g/ml. Detection: Chemiluminescence with an exposure time of 10 seconds.