Atlastin-3 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_056274.3

Catalog No. A303-312A GeneID 25923

Lot No. A303-312A-1

APPLICATIONS WB, IP

SPECIES REACTIVITY Human, Mouse

PRESUMED REACTIVITY Based on 100% sequence identity, this antibody is predicted to react with Rat

AMOUNT 100 μl

CONCENTRATION 200 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION Antibody was affinity purified using an epitope specific to Atlastin-3 immobilized on solid

PROCEDURES support.

The epitope recognized by A303-312A maps to a region between residue 350 and 400 of human

Atlastin GTPase 3 using the numbering given in entry NP_056274.3 (GeneID 25923).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 - 1:10,000

Immunoprecipitation 2 – 10 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100-020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-20% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-20% SDS-PAGE.

ADDITIONAL INFO https://www.bethyl.com/product/A303-312A


Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019





Detection of human and mouse Atlastin–3 by western blot (h and m) and immunoprecipitation (h). Samples: Whole cell lysate from HeLa (15 and 50 μg for WB; 1 mg for IP, 20% of IP loaded), Jurkat (J; 50 μg), HEK293T (T; 50 μg) and mouse NIH 3T3 (M; 50 μg) cells. Antibodies: Affinity purified rabbit anti–Atlastin–3 antibody A303–312A used for WB at 0.04 $\mu g/ml$ (A) and 1 $\mu g/ml$ (B) and used for IP at 6 $\mu g/mg$ lysate. Atlastin–3 was also immunoprecipitated by rabbit anti–Atlastin–3 antibody A303–313A, which recognizes a downstream epitope. Detection: Chemiluminescence with exposure times of 30 seconds (A) and 10 seconds (B).