Cyclin F Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_001752.2

Catalog No. A303-406A GenelD 899

Lot No. A303-406A-1

APPLICATIONS WB

SPECIES REACTIVITY Human 100 μl

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to Cyclin F immobilized on solid support.

The epitope recognized by A303-406A maps to a region between residue 736 and 786 of human

Cyclin F using the numbering given in entry NP_001752.2 (GeneID 899).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 – 1:10,000 Immunoprecipitation Not recommended

APPLICATION NOTES Western blot of lysates performed using standard western blot reagents and 4–8% SDS-PAGE.


ADDITIONAL INFO https://www.bethyl.com/product/A303-406A

Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019





Detection of human Cyclin F by western blot. Samples: Whole cell lysate from HeLa (15 and 50 μ g), HEK293T (T; 50 μ g) and Jurkat (J; 50 μ g) cells. Antibody: Affinity purified rabbit anti-Cyclin F antibody A303-406A used for WB at 0.1 μ g/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.