
INF2 Antibody						
Rabbit Polyclonal						
Antigen Affinity Purified			Protein ID	NP_071934.3		
Catalog No. A303-428A		428A	GenelD	64423	DET	
Lot No.	A303-	428A-1				ORIES, INC
APPLICATIONS		WB, IP				
SPECIES REACTIVITY		Human				
AMOUNT		100 µl				
CONCENTRATION		200 μg/ml				
STORAGE/SHELF LIFE		2 - 8° C / 1 year from date of receipt				
PHYSICAL STATE		Liquid				
BUFFER		Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide				
ISOTYPE		IgG				
ORIGIN		USA				
PRODUCTION PROCEDURES		Antibody was affinity purified using an epitope specific to INF2 immobilized on solid support.				
		The epitope recognized by A303-428A maps to a region between residue 950 and 1000 of human Inverted Formin-2 using the numbering given in entry NP_071934.3 (GeneID 64423).				
		Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.				
APPLICATIONS		Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.				
		Western Blot	1:2	2,000 - 1:10,000		
		Immunoprecip	itation 2 -	10 µg/mg lysate		
APPLICATION N	IOTES	Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).				
		Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.				
ADDITIONAL INFO		https://www.bethyl.com/product/A303-428A				
		Use the link above to view SDS, a current list of citations, and other product specific information. IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB				

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019

For in vitro laboratory use only. Not for any clinical, therapeutic or diagnostic use in humans or animals. Not for human or animal consumption. This product may not be resold or modified for resale without the prior written approval of Bethyl Laboratories, Inc. The information provided in this data sheet is believed to be correct but does not purport to be all-inclusive and is intended to be used as a guide. Bethyl Laboratories, Inc. shall not be liable or responsible in any way for use of either this information or the material supplied. Disposal of hazardous material may be subject to federal, state or local laws or regulations.



Detection of human INF2 by western blot and immunoprecipitation. *Samples:* Whole cell lysate from HeLa (15 and 50 µg for WB; 1 mg for IP, 20% of IP loaded) and HEK293T (T; 50 µg) cells. *Antibodies:* Affinity purified rabbit anti–INF2 antibody A303–428A used for WB at 0.04 µg/ml (A) and 0.4 µg/ml (B) and used for IP at 6 µg/mg lysate. INF2 was also immunoprecipitated by rabbit anti– INF2 antibodies A303–427A and A303–429A, which recognize other epitopes. *Detection:* Chemiluminescence with exposure times of 30 seconds (A) and 3 seconds (B).