DENTT Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_071400.1

Catalog No. A304-012A GeneID 64061

Lot No. A304-012A-1

APPLICATIONS WB, IP, IHC
SPECIES REACTIVITY Human

AMOUNT 100 μl

CONCENTRATION 1000 µg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to DENTT immobilized on solid support.

The epitope recognized by A304-012A maps to a region between residue 1 to 50 of human

TSPY-Like 2 using the numbering given in entry NP_071400.1 (GeneID 64061).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1,000 - 1:5,000

Immunoprecipitation $2 - 10 \mu g/mg$ lysate

Immunohistochemistry 1:500 - 1:2,000. Epitope retrieval with citrate buffer pH 6.0 is

recommended for FFPE tissue sections.

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100–020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.

IHC HUMAN CONTROLS Testis

ADDITIONAL INFO https://www.bethyl.com/product/A304-012A

Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer


Date: June 21, 2019




DENTT Antibody A304-012A




Detection of human DENTT by western blot. Samples: Whole cell lysate (50 µg) from HeLa, HEK293T, Jurkat, mouse TCMK-1, and mouse NIH 3T3 cells. Antibodies: Affinity purified rabbit anti-DENTT antibody A304-012A (lot A304-012A-1) used for WB at 0.4 µg/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.



Detection of human DENTT by western blot of immunoprecipitates. Samples: Whole cell lysate (1 mg for IP; 20% of IP loaded) from HeLa cells. Antibodies: Affinity purified rabbit anti–DENTT antibody A304–012A (lot A304–012A–1) used for IP at 6 μ g/mg lysate. DENTT was inefficiently immunoprecipitated by rabbit anti–DENTT antibody A304–013A. For blotting immunoprecipitated DENTT, A304–012A was used at 1 μ g/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.



Detection of human DENTT by immunohistochemistry. Samples: FFPE serial sections of human testis. Antibody: Affinity purified rabbit anti- DENTT (Cat. No. A304-012A Lot1, left image and Cat. No. A304-013A Lot1, right image) used at a dilution of 1:1,000 (1µg/ml). Detection: DAB



Detection of human DENTT by immunohistochemistry. Samples: FFPE serial sections of human lung cancer. Antibody: Affinity purified rabbit anti- DENTT (Cat. No. A304-012A Lot1, left image and Cat. No. A304-013A Lot1, right image) used at a dilution of 1:1,000 (1µg/ml). Detection: DAB