

ASCC3 Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID NP_006819.2

Catalog No. A304-014A


GeneID 10973

Lot No. A304-014A-1




APPLICATIONS	WB
SPECIES REACTIVITY	Human, Mouse
AMOUNT	100 µl
CONCENTRATION	1000 µg/ml
STORAGE/SHELF LIFE	2 – 8° C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antibody was affinity purified using an epitope specific to ASCC3 immobilized on solid support. The epitope recognized by A304-014A maps to a region between residue 175 to 225 of human Activating Signal Cointegrator 1 Complex Subunit 3 using the numbering given in entry NP_006819.2 (GeneID 10973). Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.
APPLICATIONS	Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot 1:2,000 – 1:10,000 Immunoprecipitation Not recommended
APPLICATION NOTES	Western blot of lysates performed using standard western blot reagents and 4–8% SDS-PAGE.
ADDITIONAL INFO	https://www.bethyl.com/product/A304-014A Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019



Detection of human and mouse ASCC3 by western blot.
Samples: Whole cell lysate (50 µg) from HEK293T, HeLa, Jurkat, mouse TCMK-1, and mouse NIH 3T3 cells.
Antibodies: Affinity purified rabbit anti-ASCC3 antibody A304-014A (lot A304-014A-1) used for WB at 0.1 µg/ml.
Detection: Chemiluminescence with an exposure time of 30 seconds.



Detection of human ASCC3 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1 mg for IP; 20% of IP loaded) from HEK293T cells. *Antibodies:* Affinity purified rabbit anti-ASCC3 antibody A304-014A (lot A304-014A-1) used for IP at 6 µg/mg lysate. ASCC3 was also immunoprecipitated by rabbit anti-ASCC3 antibody A304-015A. For blotting immunoprecipitated ASCC3, A304-014A was used at 1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 30 seconds.