GOLGB1 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_004478.3

Catalog No. A304-033A GeneID 2804

Lot No. A304-033A-1

APPLICATIONS WB, IP

SPECIES REACTIVITY Human

AMOUNT 100 μl

 $\textbf{CONCENTRATION} \qquad \quad 1000 \; \mu \text{g/ml}$

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to GOLGB1 immobilized on solid support.

The epitope recognized by A304-033A maps to a region between residue 425 to 475 of human

Golgin B1 using the numbering given in entry NP_004478.3 (GeneID 2804).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 - 1:10,000 Immunoprecipitation 2 - 10 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100–020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 3-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 3-8% SDS-PAGE.

ADDITIONAL INFO https://www.bethyl.com/product/A304-033A


Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol IP WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019







Detection of human GOLGB1 by western blot. Samples: Whole cell lysate ($50 \mu g$) from HEK293T, HeLa, and Jurkat cells. Antibodies: Affinity purified rabbit anti-GOLGB1 antibody A304-033A (lot A304-033A-1) used for WB at 0.1 $\mu g/ml$. Detection: Chemiluminescence with an exposure time of 10 seconds.

Detection of human GOLGB1 by western blot of immunoprecipitates. Samples: Whole cell lysate (1 mg for IP; 20% of IP loaded) from HEK293T cells. Antibodies: Affinity purified rabbit anti–GOLGB1 antibody A304–033A (lot A304–033A–1) used for IP at 6 μ g/mg lysate. GOLGB1 was also immunoprecipitated by rabbit anti–GOLGB1 antibodies A304–034A and A304–035A. For blotting immunoprecipitated GOLGB1, A304–033A was used at 1 μ g/ml. Detection: Chemiluminescence with an exposure time of 10 seconds.