ZNF503 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP 116161.2

Catalog No. A304-061A GeneID 84858

Lot No. A304-061A-1

APPLICATIONS WB. IP SPECIES REACTIVITY Human **AMOUNT** 100 ul

CONCENTRATION $1000 \mu g/ml$

STORAGE/SHELF LIFE 2 - 8° C / 1 year from date of receipt

PHYSICAL STATE

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG **ORIGIN USA**

PRODUCTION Antibody was affinity purified using an epitope specific to ZNF503 immobilized on solid

PROCEDURES support. The epitope recognized by A304-061A maps to a region between residue 275 to 325 of human Zinc Finger Protein 503 using the numbering given in entry NP_116161.2 (GenelD 84858).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 - 1:10,000 **Immunoprecipitation** 2 - 10 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100-020).

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.

ADDITIONAL INFO https://www.bethyl.com/product/A304-061A


Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories. Inc. Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019



ZNF503 Antibody A304-061A





Detection of human ZNF503 by western blot. Samples: Whole cell lysate (50 μg) from 293T cells. Antibodies: Affinity purified rabbit anti–ZNF503 antibody A304–061A (lot A304–061A–1) used for WB at 0.1 μg/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.

Detection of human ZNF503 by western blot of immunoprecipitates. Samples: Whole cell lysate (1 mg for IP; 20% of IP loaded) from 293T cells. Antibodies: Affinity purified rabbit anti–ZNF503 antibody A304–061A (lot A304–061A–1) used for IP at 6 μg/mg lysate. ZNF503 was also immunoprecipitated by rabbit anti–ZNF503 antibodies A304–059A and A304–060A. For blotting immunoprecipitated ZNF503, A304–061A was used at 1 μg/ml. Detection: Chemiluminescence with an exposure time of 3 seconds.