## RNF216 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP\_996999.1

Catalog No. A304-110A GeneID 54476

Lot No. A304-110A-1

APPLICATIONS IP

SPECIES REACTIVITY Human
AMOUNT 100 µl

CONCENTRATION 1000 μg/ml

**STORAGE/SHELF LIFE** 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to RNF216 immobilized on solid support.

The epitope recognized by A304-110A maps to a region between residue 1 to 50 of human Ring

Finger Protein 216 using the numbering given in entry NP\_996999.1 (GeneID 54476).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot Not recommended. Use rabbit anti-RNF216 antibody A304-111A.

Immunoprecipitation 2 – 10 µg/mg lysate

**APPLICATION NOTES** Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100–020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

ADDITIONAL INFO https://www.bethyl.com/product/A304-110A


Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol\_IP\_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019





Detection of human RNF216 by western blot of immunoprecipitates. Samples: Whole cell lysate (1 mg for IP; 20% of IP loaded) from HeLa cells. Antibodies: Affinity purified rabbit anti–RNF216 antibody A304–110A (lot A304–110A–1) used for IP at 6 μg/mg lysate. RNF216 was also immunoprecipitated by rabbit anti–RNF216 antibodies A304–111A, A304–112A and A304–113A. For blotting immunoprecipitated RNF216, A304–111A was used at 1 μg/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.