GRAMD1A Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_065946.2

Catalog No. A304-159A GenelD 57655

Lot No. A304-159A-1

APPLICATIONS IP

SPECIES REACTIVITY Human
AMOUNT 100 μI

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION Antibody was affinity purified using an epitope specific to GRAMD1A immobilized on solid

PROCEDURES support.

The epitope recognized by A304-159A maps to a region between residue 300 to 350 of human GRAM Domain Containing 1A using the numbering given in entry NP_065946.2 (GeneID 57655).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot Not recommended. Use rabbit anti-GRAMD1A antibody A304-160A.

Immunoprecipitation 2 – 10 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

ADDITIONAL INFO https://www.bethyl.com/product/A304-159A


Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019





Detection of human GRAMD1A by western blot of immunoprecipitates. Samples: Whole cell lysate (1 mg for IP; 20% of IP loaded) from HeLa cells. Antibodies: Affinity purified rabbit anti–GRAMD1A antibody A304–159A (lot A304–159A–1) used for IP at 6 μ g/mg lysate. GRAMD1A was also immunoprecipitated by rabbit anti–GRAMD1A antibodies A304–158A and A304–160A. For blotting immunoprecipitated GRAMD1A, A304–160A was used at 1 μ g/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.