FTSJ3 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_060117.3

Catalog No. A304-199A GenelD 117246

Lot No. A304-199A-1

APPLICATIONS WB, IP

SPECIES REACTIVITY Human, Mouse

PRESUMED REACTIVITY Based on 100% sequence identity, this antibody is predicted to react with Orangutan

AMOUNT 100 μl

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to FTSJ3 immobilized on solid support.

The epitope recognized by A304-199A maps to a region between residue 625 to 675 of human

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

FtsJ Homolog 3 using the numbering given in entry NP_060117.3 (GeneID 117246).

equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:10,000 - 1:25,000

Immunoprecipitation 2 – 10 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100-020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.

ADDITIONAL INFO https://www.bethyl.com/product/A304-199A

Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB


This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019



FTSJ3 Antibody A304-199A





Detection of human and mouse FTSJ3 by western blot. Samples: Whole cell lysate (50 μg) from HeLa, HEK293T, Jurkat, mouse TCMK-1, and mouse NIH 3T3 cells. Antibodies: Affinity purified rabbit anti-FTSJ3 antibody A304-199A (lot A304-199A-1) used for WB at 0.04 μg/ml. Detection: Chemiluminescence with an exposure time of 10 seconds.

Detection of human FTSJ3 by western blot of immunoprecipitates. Samples: Whole cell lysate (1 mg for IP; 20% of IP loaded) from 293T cells. Antibodies: Affinity purified rabbit anti–FTSJ3 antibody A304–199A (lot A304–199A–1) used for IP at 6 μ g/mg lysate. FTSJ3 was also immunoprecipitated by rabbit anti–FTSJ3 antibodies A304–198A and A304–200A. For blotting immunoprecipitated FTSJ3, A304–199A was used at 1 μ g/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.