TRMT1 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_060192.1

Catalog No. A304-204A GeneID 55621

Lot No. A304-204A-1

APPLICATIONS IP

AMOUNT Human 100 µl

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to TRMT1 immobilized on solid support.

The epitope recognized by A304–204A maps to a region between residue 500 to 550 of human TRM1 tRNA Methyltransferase 1 Homolog using the numbering given in entry NP_060192.1

(GeneID 55621).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot Not recommended. Use rabbit anti-TRMT1 antibody A304-205A.

Immunoprecipitation 2 – 10 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

ADDITIONAL INFO https://www.bethyl.com/product/A304-204A


Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019





Detection of human TRMT1 by western blot of immunoprecipitates. Samples: Whole cell lysate (1 mg for IP; 20% of IP loaded) from 293T cells. Antibodies: Affinity purified rabbit anti-TRMT1 antibody A304-204A (lot A304-204A-1) used for IP at 6 μg/mg lysate. TRMT1 was also immunoprecipitated by rabbit anti-TRMT1 antibody A304-205A. For blotting immunoprecipitated TRMT1, A304-205A was used at 1 μg/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.