

# RPN2 Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID NP\_002942.2

Catalog No. A304-265A

GeneID 6185

Lot No. A304-265A-1



<b>APPLICATIONS</b>	WB, IP, IHC
<b>SPECIES REACTIVITY</b>	Human
<b>PRESUMED REACTIVITY</b>	Based on 100% sequence identity, this antibody is predicted to react with Mouse, Rat, Bovine, Pig and Orangutan
<b>AMOUNT</b>	100 µl
<b>CONCENTRATION</b>	1000 µg/ml
<b>STORAGE/SHELF LIFE</b>	2 – 8° C / 1 year from date of receipt
<b>PHYSICAL STATE</b>	Liquid
<b>BUFFER</b>	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
<b>ISOTYPE</b>	IgG
<b>ORIGIN</b>	USA
<b>PRODUCTION PROCEDURES</b>	Antibody was affinity purified using an epitope specific to RPN2 immobilized on solid support.

The epitope recognized by A304-265A maps to a region between residue 350 to 400 of human Ribophorin II using the numbering given in entry NP\_002942.2 (GeneID 6185).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot	1:1,000 – 1:5,000
Immunoprecipitation	2 – 10 µg/mg lysate
Immunohistochemistry	1:500 – 1:2,000. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections.


**APPLICATION NOTES** Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.


**IHC HUMAN CONTROLS** Prostate Carcinoma

**ADDITIONAL INFO** <https://www.bethyl.com/product/A304-265A>  
Use the link above to view SDS, a current list of citations, and other product specific information.  
IP-western blot protocol: [https://www.bethyl.com/content/protocol\\_IP\\_WB](https://www.bethyl.com/content/protocol_IP_WB)


This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.  
Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019




**Detection of human RPN2 by western blot.** *Samples:* Whole cell lysate (50 µg) from HeLa, HEK293T, and Jurkat cells. *Antibodies:* Affinity purified rabbit anti-RPN2 antibody A304-265A (lot A304-265A-1) used for WB at 0.4 µg/ml. *Detection:* Chemiluminescence with an exposure time of 3 minutes.



**Detection of human RPN2 by western blot of immunoprecipitates.** *Samples:* Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from HeLa cells. *Antibodies:* Affinity purified rabbit anti-RPN2 antibody A304-265A (lot A304-265A-1) used for IP at 6 µg per reaction. RPN2 was also immunoprecipitated by rabbit anti-RPN2 antibody BL15802. For blotting immunoprecipitated RPN2, A304-265A was used at 1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 30 seconds.



**Detection of human RPN2 by immunohistochemistry.** *Sample:* FFPE section of human mammary glands. *Antibody:* Affinity purified rabbit anti-RPN2 (Cat. No. A304-265A Lot1) used at a dilution of 1:1,000 (1 µg/ml). *Detection:* DAB



**Detection of human RPN2 by immunohistochemistry.** *Sample:* FFPE section of human prostate carcinoma. *Antibody:* Affinity purified rabbit anti-RPN2 (Cat. No. A304-265A Lot1) used at a dilution of 1:1,000 (1 µg/ml). *Detection:* DAB