Syntaxin 6 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_005810.1

Catalog No. A304-397A GeneID 10228

Lot No. A304-397A-1

APPLICATIONS WB, IP

SPECIES REACTIVITY Human, Mouse

PRESUMED REACTIVITY Based on 100% sequence identity, this antibody is predicted to react with Orangutan

AMOUNT 100 μl

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION Antibody was affinity purified using an epitope specific to Syntaxin 6 immobilized on solid

PROCEDURES support.

The epitope recognized by A304-397A maps to a region between residue 75 to 125 of human

Syntaxin 6 using the numbering given in entry NP_005810.1 (GeneID 10228).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 - 1:10,000

Immunoprecipitation 2 – 10 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100-020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-20% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-20% SDS-PAGE.

ADDITIONAL INFO https://www.bethyl.com/product/A304-397A


Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019







Detection of human and mouse Syntaxin 6 by western blot. Samples: Whole cell lysate (50 μg) prepared using NETN or RIPA buffer from HeLa, HEK293T, Jurkat, mouse TCMK-1, and mouse NIH 3T3 cells. Antibodies: Affinity purified rabbit anti-Syntaxin 6 antibody A304-397A (lot A304-397A-1) used for WB at 0.1 μg/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.

Detection of human Syntaxin 6 by western blot of immunoprecipitates. Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) prepared using RIPA buffer from Jurkat cells. Antibodies: Affinity purified rabbit anti–Syntaxin 6 antibody A304–397A (lot A304–397A–1) used for IP at 6 μg per reaction. Syntaxin 6 was also immunoprecipitated by rabbit anti–Syntaxin 6 antibody BL16441. For blotting immunoprecipitated Syntaxin 6, A304–397A was used at 1 μg/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.