ARHGAP5 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_001025226.1

Catalog No. A304-548A GeneID 394

Lot No. A304-548A-1

APPLICATIONS WB, IP

SPECIES REACTIVITY Human, Mouse

AMOUNT 100 μl

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION Antibody was affinity purified using an epitope specific to ARHGAP5 immobilized on solid

PROCEDURES support.

The epitope recognized by A304–548A maps to a region between residue 1050 to 1100 of

human Rho GTPase Activating Protein 5 using the numbering given in entry NP_001025226.1

(GeneID 394).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 - 1:10,000

Immunoprecipitation 2 – 10 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100-020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.

ADDITIONAL INFO https://www.bethyl.com/product/A304-548A


Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019







Detection of human and mouse ARHGAP5 by western blot. Samples: Whole cell lysate (50 μg) from HeLa, HEK293T, Jurkat, mouse TCMK-1, and mouse NIH 3T3 cells prepared using NETN lysis buffer. Antibodies: Affinity purified rabbit anti-ARHGAP5 antibody A304-548A (lot A304-548A-1) used for WB at 0.1 μg/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.

Detection of human ARHGAP5 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from 293T cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-ARHGAP5 antibody A304-548A (lot A304-548A-1) used for IP at 6 μg per reaction. ARHGAP5 was also immunoprecipitated by rabbit anti-ARHGAP5 antibody A304-549A. For blotting immunoprecipitated ARHGAP5, A304-548A was used at 1 μg/ml. *Detection:* Chemiluminescence with an exposure time of 3 minutes.