UHRF1BP1 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_060224.3

Catalog No. A304-646A GeneID 54887

Lot No. A304-646A-1

APPLICATIONS WB, IP

SPECIES REACTIVITY Human

AMOUNT 100 μl

CONCENTRATION 1000 μg/ml

STORAGE/SHELF LIFE 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION Antibody was affinity purified using an epitope specific to UHRF1BP1 immobilized on solid

PROCEDURES support.

The epitope recognized by A304-646A maps to a region between residue 1390 to 1440 of human UHRF1 Binding Protein 1 using the numbering given in entry NP 060224.3 (GeneID

54887).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1,000 - 1:5,000

Immunoprecipitation 2 – 10 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100-020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-8% SDS-PAGE.


ADDITIONAL INFO https://www.bethyl.com/product/A304-646A


Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019







Detection of human UHRF1BP1 by western blot. Samples: Whole cell lysate (50 μg) from HeLa, HEK293T, and Jurkat cells prepared using NETN lysis buffer. Antibodies: Affinity purified rabbit anti-UHRF1BP1 antibody A304-646A (lot A304-646A-1) used for WB at 0.4 μg/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.

Detection of human UHRF1BP1 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from 293T cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-UHRF1BP1 antibody A304-646A (lot A304-646A-1) used for IP at 6 μg per reaction. UHRF1BP1 was also immunoprecipitated by rabbit anti-UHRF1BP1 antibody BL17846. For blotting immunoprecipitated UHRF1BP1, A304-646A was used at 1 μg/ml. *Detection:* Chemiluminescence with an exposure time of 30 seconds.