## TBC1D9B Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP\_942568.2

Catalog No. A304-654A GeneID 23061

Lot No. A304-654A-1

APPLICATIONS IP

SPECIES REACTIVITY Human
AMOUNT 100 µI

CONCENTRATION 1000 μg/ml

**STORAGE/SHELF LIFE** 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

**PRODUCTION** Antibody was affinity purified using an epitope specific to TBC1D9B immobilized on solid

**PROCEDURES** support.

The epitope recognized by A304-654A maps to a region between residue 1050 to 1100 of human TBC1 Domain Family. Member 9B using the numbering given in entry NP 942568.2

(GeneID 23061).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot Not recommended. Use rabbit anti-TBC1D9B antibody A304-655A.

Immunoprecipitation 2 – 10 μg/mg lysate

**APPLICATION NOTES** Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100-020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE

(link to IP-western blot protocol in Additional Info section below).


ADDITIONAL INFO https://www.bethyl.com/product/A304-654A

Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol IP WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019





Detection of human TBC1D9B by western blot of immunoprecipitates. Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from 293T cells prepared using NETN lysis buffer. Antibodies: Affinity purified rabbit anti-TBC1D9B antibody A304-654A (lot A304-654A-1) used for IP at 6 μg per reaction. TBC1D9B was also immunoprecipitated by rabbit anti-TBC1D9B antibody A304-655A. For blotting immunoprecipitated TBC1D9B, A304-654A was used at 1 μg/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.