

EIF3M Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID Q7L2H7.1

Catalog No. A304-761A

GeneID 10480

Lot No. A304-761A-1



APPLICATIONS	WB, IP
SPECIES REACTIVITY	Human, Mouse
PRESUMED REACTIVITY	Based on 100% sequence identity, this antibody is predicted to react with Bovine and Orangutan
AMOUNT	100 µl
CONCENTRATION	1000 µg/ml
STORAGE/SHELF LIFE	2 - 8° C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antibody was affinity purified using an epitope specific to EIF3M immobilized on solid support.

The epitope recognized by A304-761A maps to a region between residue 1 to 50 of human Eukaryotic Translation Initiation Factor 3 Subunit M using the numbering given in entry Q7L2H7.1 (GeneID 10480).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 - 1:10,000

Immunoprecipitation 2 - 10 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-20% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-20% SDS-PAGE.

ADDITIONAL INFO


<https://www.bethyl.com/product/A304-761A>

Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB


This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019



Detection of human and mouse EIF3M by western blot.

Samples: Whole cell lysate (50 µg) from HeLa, HEK293T, Jurkat, mouse TCMK-1, and mouse NIH 3T3 cells prepared using NETN lysis buffer. **Antibodies:** Affinity purified rabbit anti-EIF3M antibody A304-761A (lot A304-761A-1) used for WB at 0.1 µg/ml. **Detection:** Chemiluminescence with an exposure time of 1 second.



Detection of human EIF3M by western blot of immunoprecipitates.

Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from 293T cells prepared using NETN lysis buffer. **Antibodies:** Affinity purified rabbit anti-EIF3M antibody A304-761A (lot A304-761A-1) used for IP at 6 µg per reaction. EIF3M was also immunoprecipitated by rabbit anti-EIF3M antibody BL18577. For blotting immunoprecipitated EIF3M, A304-761A was used at 0.4 µg/ml. **Detection:** Chemiluminescence with an exposure time of 1 second.