

# Mdc1 /Nfbd1 Antibody


Rabbit Polyclonal

Antigen Affinity Purified	Protein ID	Q5PSV9
Catalog No. A304-843A	GeneID	240087
Lot No. A304-843A-1		



<b>APPLICATIONS</b>	IP
<b>SPECIES REACTIVITY</b>	Mouse
<b>AMOUNT</b>	100 µl
<b>CONCENTRATION</b>	1000 µg/ml
<b>STORAGE/SHELF LIFE</b>	2 – 8° C / 1 year from date of receipt
<b>PHYSICAL STATE</b>	Liquid
<b>BUFFER</b>	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
<b>ISOTYPE</b>	IgG
<b>ORIGIN</b>	USA
<b>PRODUCTION PROCEDURES</b>	<p>Antibody was affinity purified using an epitope specific to Mdc1 /Nfbd1 immobilized on solid support.</p> <p>The epitope recognized by A304-843A maps to a region between residue 900 to 950 of human Mdc1 /Nfbd1 using the numbering given in entry Q5PSV9 (GeneID 240087).</p> <p>Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.</p>
<b>APPLICATIONS</b>	<p>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.</p> <p>Western Blot                      Not recommended. A304-843A has not performed satisfactorily when used for WB of Mdc1 /Nfbd1 in crude preparations (e.g. whole cell lysate). This antibody can be used for WB of enriched (e.g. immunoprecipitated) sources of Mdc1 /Nfbd1.</p> <p>Immunoprecipitation      2 – 10 µg/mg lysate</p>
<b>APPLICATION NOTES</b>	Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).
<b>ADDITIONAL INFO</b>	<p><a href="https://www.bethyl.com/product/A304-843A">https://www.bethyl.com/product/A304-843A</a></p> <p>Use the link above to view SDS, a current list of citations, and other product specific information. IP-western blot protocol: <a href="https://www.bethyl.com/content/protocol_IP_WB">https://www.bethyl.com/content/protocol_IP_WB</a></p>

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.  
Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019



**Detection of mouse Mdc1/Nfbd1 by western blot of immunoprecipitates.** *Samples:* Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from TCMK-1 cells prepared using NETN lysis buffer. *Antibody:* Affinity purified rabbit anti-Mdc1/Nfbd1 antibody A304-843A (lot A304-843A-1) used for IP at 6 µg per reaction. Mdc1/Nfbd1 was also immunoprecipitated by rabbit anti-Mdc1/Nfbd1 antibody BL18630. For blotting immunoprecipitated Mdc1/Nfbd1, A304-843A was used at 1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 3 minutes.