## **RABGAP1** Antibody

Rabbit Polyclonal

Catalog No.

Antigen Affinity Purified Protein ID Q9Y3P9.3

Lot No. A304-913A-1

A304-913A

APPLICATIONS IP

SPECIES REACTIVITY Human

PRESUMED REACTIVITY Based on 100% sequence identity, this antibody is predicted to react with Mouse and Orangutan

23637

AMOUNT 100 μl

CONCENTRATION 1000 μg/ml

**STORAGE/SHELF LIFE** 2 – 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

**BUFFER** Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

GeneID

ISOTYPE IgG
ORIGIN USA

**PRODUCTION** Antibody was affinity purified using an epitope specific to RABGAP1 immobilized on solid

**PROCEDURES** support.

The epitope recognized by A304-913A maps to a region between residue 25 to 75 of human Rab

GTPase-activating protein 1 using the numbering given in entry Q9Y3P9.3 (GeneID 23637).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4

equals 1.0 mg of IgG.

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot Not recommended, Use rabbit anti-RABGAP1 antibody A304-912A.

Immunoprecipitation 2 – 10 µg/mg lysate

**APPLICATION NOTES** Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. \$100-020),

Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE


(link to IP-western blot protocol in Additional Info section below).


ADDITIONAL INFO https://www.bethyl.com/product/A304-913A

Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol\_IP\_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019





Detection of human RABGAP1 by western blot of immunoprecipitates. Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from HeLa cells prepared using NETN lysis buffer. Antibodies: Affinity purified rabbit anti-RABGAP1 antibody A304–913A (lot A304–913A–1) used for IP at 6 μg per reaction. RABGAP1 was also immunoprecipitated by rabbit anti-RABGAP1 antibody BL19305. For blotting immunoprecipitated RABGAP1, A304–913A was used at 1 μg/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.