

SPATS2/SPATA10 Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID Q86XZ4.1

Catalog No. A304-970A

GeneID 65244

Lot No. A304-970A-1



APPLICATIONS WB, IP

SPECIES REACTIVITY Human

AMOUNT 100 µl

CONCENTRATION 1000 µg/ml

STORAGE/SHELF LIFE 2 - 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG

ORIGIN USA

PRODUCTION PROCEDURES Antibody was affinity purified using an epitope specific to SPATS2/SPATA10 immobilized on solid support.

The epitope recognized by A304-970A maps to a region between residue 450 to 500 of human Spermatogenesis-associated serine-rich protein 2 using the numbering given in entry Q86XZ4.1 (GeneID 65244).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 - 1:10,000

Immunoprecipitation 2 - 10 µg/mg lysate

APPLICATION NOTES Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-20% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).

Western blot of lysates performed using standard western blot reagents and 4-20% SDS-PAGE.


ADDITIONAL INFO <https://www.bethyl.com/product/A304-970A>

Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019



Detection of human SPATS2 by western blot. *Samples:* Whole cell lysate (10 µg) from HeLa, HEK293T, and Jurkat cells prepared using NETN lysis buffer. *Antibody:* Affinity purified rabbit anti-SPATS2 antibody A304-970A (lot A304-970A-1) used for WB at 0.1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 3 minutes.



Detection of human SPATS2 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from HeLa cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-SPATS2 antibody A304-970A (lot A304-970A-1) used for IP at 6 µg per reaction. SPATS2 was also immunoprecipitated by rabbit anti-SPATS2 antibody A304-971A. For blotting immunoprecipitated SPATS2, A304-970A was used at 1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 3 minutes.