

GGA1 Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID Q9UJY5.1

Catalog No. A305-320A

GeneID 26088


Lot No. A305-320A-1



APPLICATIONS	IP				
SPECIES REACTIVITY	Human				
AMOUNT	100 µl				
CONCENTRATION	1000 µg/ml				
STORAGE/SHELF LIFE	2 – 8° C / 1 year from date of receipt				
PHYSICAL STATE	Liquid				
BUFFER	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide				
ISOTYPE	IgG				
ORIGIN	USA				
PRODUCTION PROCEDURES	Antibody was affinity purified using an epitope specific to GGA1 immobilized on solid support. The epitope recognized by A305-320A maps to a region between residue 300 to 350 of human ADP-ribosylation factor-binding protein GGA1 using the numbering given in entry Q9UJY5.1 (GeneID 26088). Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.				
APPLICATIONS	Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. <table><tr><td>Western Blot</td><td>Not recommended. A305-320A has not performed satisfactorily when used for WB of GGA1 in crude preparations (e.g. whole cell lysate). This antibody can be used for WB of enriched (e.g. immunoprecipitated) sources of GGA1.</td></tr><tr><td>Immunoprecipitation</td><td>2 – 10 µg/mg lysate</td></tr></table>	Western Blot	Not recommended. A305-320A has not performed satisfactorily when used for WB of GGA1 in crude preparations (e.g. whole cell lysate). This antibody can be used for WB of enriched (e.g. immunoprecipitated) sources of GGA1.	Immunoprecipitation	2 – 10 µg/mg lysate
Western Blot	Not recommended. A305-320A has not performed satisfactorily when used for WB of GGA1 in crude preparations (e.g. whole cell lysate). This antibody can be used for WB of enriched (e.g. immunoprecipitated) sources of GGA1.				
Immunoprecipitation	2 – 10 µg/mg lysate				
APPLICATION NOTES	Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-8% SDS-PAGE (link to IP-western blot protocol in Additional Info section below).				
ADDITIONAL INFO	https://www.bethyl.com/product/A305-320A Use the link above to view SDS, a current list of citations, and other product specific information. IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB				

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019



Detection of human GGA1 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from 293T cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-GGA1 antibody A305-320A (lot A305-320A-1) used for IP at 6 µg per reaction. GGA1 was also immunoprecipitated by rabbit anti-GGA1 antibody A305-368A. For blotting immunoprecipitated GGA1, A305-320A was used at 0.4 µg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.