

Goat IgG-heavy and light chain cross-adsorbed Antibody

F(ab')₂ Donkey Polyclonal

Antigen Affinity Purified

Catalog No. A50-207A

Lot No. A50-207A-4



APPLICATIONS	WB, IHC, ICC, ELISA								
SPECIES REACTIVITY	Goat. Minimum reactivity to chicken, human, mouse, rabbit and rat								
ISOTYPE	IgG								
AMOUNT	1 ml at 0.5 mg/ml								
STORAGE/SHELF LIFE	2 – 8° C / 1 year from date of receipt								
PHYSICAL STATE	Liquid								
BUFFER	Phosphate Buffered Saline (PBS) containing 0.09% Sodium Azide								
ORIGIN	USA								
PRODUCTION PROCEDURES	Antiserum was cross adsorbed using chicken, human, mouse, rabbit and rat immunosorbents to remove cross reactive antibodies. The antibody to goat IgG was isolated by affinity chromatography using antigen coupled to agarose beads. F(ab')₂ fragments were generated using a pepsin digestion. Fc fragments and whole IgG molecules have been removed. Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. By immunoelectrophoresis and ELISA this antibody reacts specifically with goat IgG and with light chains common to other goat immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. Less than 1% cross reactivity to chicken, human, mouse, rabbit and rat IgG was detected. This antibody may cross react with IgG from other species.								
APPLICATIONS	Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. <table><tr><td>Western Blot</td><td>1:1,000 – 1:20,000</td></tr><tr><td>Immunohistochemistry</td><td>1:200 – 1:2,000</td></tr><tr><td>Immunocytochemistry</td><td>1:200 – 1:2,000</td></tr><tr><td>ELISA</td><td>1:1,000 – 1:20,000; for coating plates 1:50 – 1:250</td></tr></table>	Western Blot	1:1,000 – 1:20,000	Immunohistochemistry	1:200 – 1:2,000	Immunocytochemistry	1:200 – 1:2,000	ELISA	1:1,000 – 1:20,000; for coating plates 1:50 – 1:250
Western Blot	1:1,000 – 1:20,000								
Immunohistochemistry	1:200 – 1:2,000								
Immunocytochemistry	1:200 – 1:2,000								
ELISA	1:1,000 – 1:20,000; for coating plates 1:50 – 1:250								
APPLICATION NOTES	Not all listed applications have been specifically tested by our laboratory.								
ADDITIONAL INFO	http://www.bethyl.com/product/A50-207A Use the link above to view SDS, a current list of citations, and other product specific information.								