

RelA Recombinant Monoclonal Antibody [BLR361N]

Rabbit Recombinant Monoclonal

Purified		RefSeq ID	NP_068810.3
Catalog No.	A700-361	Uniprot ID	Q04206
Lot No.	1	GenelD	5970

APPLICATIONS	WB, IP, IHC, ICC								
SPECIES REACTIVITY	Human, Mouse								
AMOUNT	100 µl (50+ tests)								
CONCENTRATION	50 µg/ml								
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt								
PHYSICAL STATE	Liquid								
BUFFER	Phosphate Buffered Saline (PBS) with 0.1% BSA and 0.09% Sodium Azide								
ISOTYPE	IgG								
CLONE #	BLR361N								
ORIGIN	USA								
PRODUCTION PROCEDURES	Recombinant antibody was purified from cell culture supernatant. Immunogen was a peptide representing a region between residue 501 and the C-terminus (residue 551) of human RelA (aka transcription factor p65) using the numbering given in entry NP_06881								
APPLICATIONS	Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. <table><tr><td>Western Blot</td><td>1:1,000</td></tr><tr><td>Immunoprecipitation</td><td>20 µl/mg lysate</td></tr><tr><td>Immunohistochemistry</td><td>1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections.</td></tr><tr><td>Immunocytochemistry</td><td>1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE cell sections.</td></tr></table>	Western Blot	1:1,000	Immunoprecipitation	20 µl/mg lysate	Immunohistochemistry	1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections.	Immunocytochemistry	1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE cell sections.
Western Blot	1:1,000								
Immunoprecipitation	20 µl/mg lysate								
Immunohistochemistry	1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE tissue sections.								
Immunocytochemistry	1:100 to 1:500. Epitope retrieval with citrate buffer pH 6.0 is recommended for FFPE cell sections.								
APPLICATION NOTES	All western blot analysis is performed using 5% Milk-TBST for blocking and as antibody diluent. Primary antibody is incubated overnight. Western blots of cell lysates are performed using Goat anti-Rabbit IgG Heavy and Light Chain Antibody (A120-101P). Western blots of immunoprecipitates are performed using Goat anti-Rabbit Light Chain HRP Conjugate (A120-113P) with 5% Normal Pig Serum (S100-020) added to the blocking buffer.								
IHC HUMAN CONTROLS	Breast Carcinoma, Testicular Seminoma, Tonsil, Cal-27 Cells, OVCAR-8 Cells								
IHC MOUSE CONTROLS	Testis, Thymus, 4T1 Cells								
ADDITIONAL INFO	https://www.fortislifesciences.com/p/A700-361 Use the link above to view SDS, a current list of citations, and other product specific information.								

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: March 28, 2025