

Mouse IgG Heavy and Light Chain Cross-Adsorbed Antibody

Rabbit Polyclonal Conjugate DyLight® 680

Antigen Affinity Purified

Catalog No. A90-317D6

Lot No. 5

APPLICATIONS	WB, IHC, ICC, Flow Cyt, IF
SPECIES REACTIVITY	Mouse. Minimum reactivity to human and rat
AMOUNT	1 ml
CONCENTRATION	0.5 mg/ml
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide
FLUOROPHORE/PROTEIN	3.5
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antiserum was cross adsorbed using human and rat immunosorbents to remove cross reactive antibodies. The antibody to mouse IgG was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to DyLight® 680.

Immunoglobulin concentration was determined using Beer's Law where 1 mg/mL IgG has an A280 of 1.4.

By immunoelectrophoresis and ELISA this antibody reacts specifically with mouse IgG and with light chains common to other mouse immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. Less than 1% cross reactivity to human and rat IgG was detected. This antibody may cross react with IgG from other species.

APPLICATIONS	Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.
	Western Blot 1:1000 – 1:20,000. 5% non-fat dry milk in PBST or TBST is recommended for blocking and incubation of antibodies. BSA is not recommended.
	Immunohistochemistry 1:50 – 1:500
	Immunocytochemistry 1:50 – 1:500
	Flow Cytometry 1:50 – 1:200
	Immunofluorescence 1:50 – 1:500


APPLICATION NOTES Not all listed applications have been specifically tested by our laboratory.

DyLight® 680 is excited at 682 (in PBS) and emits at 715 (in PBS).

DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.

ADDITIONAL INFO Please visit our website for additional product information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: August 13, 2025



Detection of GAPDH and hnRNP in HeLa Whole Cell

Lysate. *Primary Antibodies:* cocktail of goat anti-GAPDH A303-878A (A303-878A-1) and mouse anti-hnRNP A500-011A (A500-011A-1) at 1 µg/ml each. *Secondary Antibodies:* cocktail of Dylight® 800-conjugated rabbit anti-goat A50-200D8 (A50-200D8-1) (blue) and Dylight® 680-conjugated rabbit anti-mouse A90-317D6 (A90-317D6-1) (red) at 0.5 µg/ml each. *Acquisition:* Syngene G:Box, 47 seconds (green) and 25 seconds (red).