

GOAT SERUM (Lyophilized) - D304-05

Code : D304-05		Size: 5 mL	
Product Description: GOAT SERUM (Lyophilized) - D304-05			
Concentration: 95 mg/mL by Refractometry			
PhysicalState: Lyophilized			
Label	Unconjugate	Unconjugated	
Buffer	0.02 M Pota	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2	
Reconstitution Volume	5.0 mL	5.0 mL	
Reconstitution Buffer	Restore with	Restore with deionized water (or equivalent)	
Stabilizer	None	None	
Preservative	0.01% (w/v)	0.01% (w/v) Sodium Azide	
Storage Condition	Avoid cycles temperature	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.	
Synonyms	Goat serum	Goat serum for bioassay, control Goat serum, non sterile serum from Goat, non-sterile Goat serum	
Application Note		Suitable for use as a control or diluent in western blot, ELISA, immunohistochemistry assays and most other immunological methods.	
Background	macromolect low molecula Normal Goa have been s and non-fat maximizing a primary antil	Goat Serum is used as a supplement to cell culture media. Goat Serum provides a broad spectrum of macromolecules, carrier proteins for lipoid substances and trace elements, attachment and spreading factors, low molecular weight nutrients, and hormones and growth factors that promote cell growth and health. Normal Goat Serum can be used as a blocking agent to treat plastic surfaces, membrane or tissue after they have been sensitized with primary antibody or antigen. It provides an alternative to bovine serum albumin (BSA) and non-fat dry milk. It is effective in reducing nonspecific binding of proteins to reaction surfaces, thereby maximizing signal-to-noise ratio. This blocking agent is recommended for use in immunoassays where the primary antibody was produced in goat, as a source of non-specific serum protein or on tissue for immunohistochemical applications.	
Purity And Specificity		t was prepared from normal serum by delipidation and defibrination. Assay by trophoresis resulted in multiple precipitin arcs against anti-Goat Serum.	
Expiration	Expiration d	Expiration date is one (1) year from date of opening.	
Related Products			
	200-301-268	Anti-AKT pS473 (MOUSE) Monoclonal Antibody - 200-301-268	
	610-4302	Anti-MOUSE IgG (H&L) (RABBIT) Antibody Peroxidase Conjugated - 610-4302	
	611-1302	Anti-RABBIT IgG (H&L) (GOAT) Antibody Peroxidase Conjugated - 611-1302	
	B304	NORMAL GOAT SERUM (NGS) - B304	
Images			
	1	Normal Goat Serum can be used as a blocking agent to treat plastic surfaces, membrane or tissue after they have been sensitized with primary antibody or antigen. It provides an alternative to bovine serum albumin (BSA) and non-fat dry milk. It is effective in reducing nonspecific binding of proteins to reaction surfaces, thereby maximizing signal-to-noise ratio. This blocking agent is recommended for use in immunoassays where the primary antibody was produced in goat, as a source of non- specific serum protein or on tissue for immunohistochemical applications. This product was prepared from normal serum by delipidation and defibrination.	



Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.