

GOAT Gamma Globulin Fraction - D604-1000

Code: D604-1000

Size: 1 g

Product Description: GOAT Gamma Globulin Fraction - D604-1000

Concentration: 40 mg/mL by UV absorbance at 280 nm

PhysicalState: Lyophilized


Label	Unconjugated
Buffer	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reconstitution Volume	100 mL
Reconstitution Buffer	Restore with deionized water (or equivalent)
Stabilizer	None
Preservative	None
Storage Condition	Store gamma globulin at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Synonyms	Plasma Gamma Globulin, Serum Gamma Globulin, Globulin Fractions, Gammaglobulin, goat gamma fraction
Application Note	Gamma Globulin Fractions can be utilized in Western Blotting, ELISA experiments, and other detection assays.
Background	Goat gamma globulin blocking reagent is an ideal blocker for western blotting, ELISA, Immunohistochemistry and other detection assays. Gamma globulins are a class of globulins, identified by their position after serum protein electrophoresis. The most significant gamma globulins are immunoglobulins ("Igs"), more commonly known as antibodies, although some Igs are not gamma globulins, and some gamma globulins are not Igs. Globulin is one of the three main types of serum proteins, the others being albumin and fibrinogen.
Purity And Specificity	Goat Gamma was prepared from normal goat serum by a multi-step process which includes delipidation and salt fractionation followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in precipitin arcs against anti-Goat Serum corresponding to gamma globulins.
ELISA	User Optimized
Western Blot	User Optimized
Immunohistochemistry	User Optimized
IF Microscopy	User Optimized
Expiration	Expiration date is one (1) year from date of opening.

Related Products

200-301-268	Anti-AKT pS473 (MOUSE) Monoclonal Antibody - 200-301-268
610-4302	Anti-MOUSE IgG (H&L) (RABBIT) Antibody Peroxidase Conjugated - 610-4302
611-1302	Anti-RABBIT IgG (H&L) (GOAT) Antibody Peroxidase Conjugated - 611-1302
B304	NORMAL GOAT SERUM (NGS) - B304


Images

1	SDS-PAGE of Goat Gamma Globulin Fraction. Lane 1: Non-reduced Goat Gamma Globulin Fraction. M: 5µL OPAL Pre-stained Marker (MB-210-0500). Lane 2: Reduced Goat Gamma Globulin Fraction. Load: 1 µg per lane. Predicted/Observed size: Non-reduced at 150-170 kDa , Reduced at 55, 25 kDa. Non-reduced migrates slightly higher.
---	---



2

SDS-PAGE results of Goat Gamma Globulin. Lane 1: reduced Goat Gamma Globulin. Lane 2: Opal PreStained Molecular Weight Ladder (p/n MB-210-0500). Lane 3: non-reduced Goat Gamma Globulin. Load: 1.0µg. 4-20% SDS Gel, Coomassie Blue Stained. Expect: reduced at 55, 25 kDa; non-reduced 150-170kDa, migrates higher.



Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.