




Detection of human HDAC1 by immunohistochemistry.
Sample: FFPE section of human colon adenocarcinoma.
Antibody: Affinity purified rabbit anti-HDAC1 (Cat. No. IHC-00056) used at a dilution of 1:250. *Detection:* DAB



Detection of mouse HDAC1 by immunohistochemistry.
Sample: FFPE section of mouse squamous cell carcinoma.
Antibody: Affinity purified rabbit anti-HDAC1 (Cat. No. IHC-00056) used at a dilution of 1:250. *Detection:* DAB



Detection of human HDAC1 by immunohistochemistry.
Sample: FFPE section of human breast carcinoma.
Antibody: Affinity purified rabbit anti-HDAC1 (Cat. No. IHC-00056) used at a dilution of 1:250. *Detection:* Green-fluorescent Anti-rabbit IgG FITC conjugated (Cat. No. A120-501F) used at a dilution of 1:100.



Detection of human HDAC1 by immunohistochemistry.
Sample: FFPE section of human breast carcinoma.
Antibody: Affinity purified rabbit anti-HDAC1 (Cat. No. IHC-00056) used at a dilution of 1:250. *Detection:* Green-fluorescent Anti-rabbit IgG FITC conjugated (Cat. No. A120-501F) used at a dilution of 1:100.