

XRN2 IHC Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID NP_036387.2

Catalog No. IHC-00223


GeneID 22803

Lot No. IHC-00223-1




APPLICATIONS	IHC, IHC-IF
SPECIES REACTIVITY	Human, Mouse
PRESUMED REACTIVITY	Based on 100% sequence identity, this antibody is predicted to react with Orangutan, Monkey, Chimpanzee, Northern white-cheeked gibbon, Little brown bat, African elephant and Crab-eating macaque
AMOUNT	100 µl
CONCENTRATION	250 µg/ml
STORAGE/SHELF LIFE	2 – 8° C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antibody was affinity purified using an epitope specific to XRN2 immobilized on solid support. The epitope recognized by IHC-00223 maps to a region between residue 900 and 950 human 5'-3' exoribonuclease 2 using the numbering given in entry NP_036387.2 (GeneID 22803). Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.
APPLICATIONS	Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Immunohistochemistry 1:100 – 1:500 Immunofluorescence 1:50 – 1:500 (IHC)
APPLICATION NOTES	Epitope exposure is recommended. Epitope exposure with citrate buffer will enhance staining. Likely to work with frozen sections. In some cases, the antibody may be diluted further than indicated.
IHC HUMAN CONTROLS	Breast Carcinoma, Colon Carcinoma, Linitis Plastica Stomach Cancer, Non-Small Cell Lung Cancer, Ovarian Carcinoma, Pancreatic Islet Cell Tumor, Prostate Carcinoma, Skin Basal Cell Carcinoma, Small Cell Lung Cancer, Stomach Adenocarcinoma, Testicular Seminoma
IHC MOUSE CONTROLS	Colon Carcinoma CT26, Teratoma
ADDITIONAL INFO	https://www.bethyl.com/product/IHC-00223 Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019

**Detection of human XRN2 by immunohistochemistry.**


Sample: FFPE section of human breast carcinoma.

Antibody: Affinity purified rabbit anti-XRN2 (Cat. No. IHC-00223) used at a dilution of 1:250. *Detection:* DAB

**Detection of human XRN2 by immunohistochemistry.**


Sample: FFPE section of human ovarian carcinoma.

Antibody: Affinity purified rabbit anti-XRN2 (Cat. No. IHC-00223) used at a dilution of 1:250. *Detection:* DAB

**Detection of human XRN2 by immunohistochemistry.**

Sample: FFPE section of human prostate carcinoma.

Antibody: Affinity purified rabbit anti-XRN2 (Cat. No. IHC-00223) used at a dilution of 1:100. *Detection:* Red-fluorescent goat anti-rabbit IgG highly cross-adsorbed Antibody.

**Detection of human XRN2 by immunohistochemistry.**

Sample: FFPE section of human breast carcinoma.

Antibody: Affinity purified rabbit anti-XRN2 (Cat. No. IHC-00223) used at a dilution of 1:100. *Detection:* Red-fluorescent Alexa Fluor® 555 goat anti-rabbit IgG (Invitrogen) used at a dilution of 1:500.