
NRBF2 IHC Antibody


| Rabbit Polyclo | | | | | |
|---|-----------|--|------------|-------------|--------|
| Antigen Affinity Purified | | | Protein ID | NP_110386.2 | |
| Catalog No. | IHC-00359 | | GenelD | 29982 | RETHVI |
| Lot No. | IHC-00 | 359-1 | | | |
| APPLICATIONS | | IHC | | | |
| SPECIES REACTIVITY | | Human, Mouse | | | |
| PRESUMED REACTIVITY | | Based on 100% sequence identity, this antibody is predicted to react with Orangutan, Monkey, Gorilla, Chimpanzee, Northern white-cheeked gibbon and Duckbill platypus 100 µl | | | |
| CONCENTRATION | | 250 µg/ml | | | |
| | | 2 - 8° C / 1 year from date of receipt | | | |
| STORAGE/SHELF LIFE | | | | | |
| PHYSICAL STATE | | Liquid | | | |
| BUFFER | | Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide | | | |
| ISOTYPE | | IgG | | | |
| ORIGIN | | USA | | | |
| PRODUCTION PROCEDURES | | Antibody was affinity purified using an epitope specific to NRBF2 immobilized on solid support. | | | |
| PROCEDURES | | The epitope recognized by IHC-00359 maps to a region between residue 125 and 175 of human nuclear receptor binding factor 2 using the numbering given in entry NP_110386.2 (GeneID 29982). | | | |
| | | Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG. | | | |
| APPLICATIONS | | Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. | | | |
| | | Immunohistochemistry 1:100 – 1:500 | | | |
| APPLICATION NOTES | | Epitope exposure is recommended. | | | |
| | | Epitope exposure with citrate buffer will enhance staining. | | | |
| | | Likely to work with frozen sections. | | | |
| IHC HUMAN CONTROLS IHC MOUSE CONTROLS ADDITIONAL INFO | | In some cases, the antibody may be diluted further than indicated. Breast Carcinoma, Linitis Plastica Stomach Cancer, Ovarian Carcinoma, Prostate Carcinoma, Testicular Seminoma | | | |
| | | Renal Cell Carcinoma, Teratoma | | | |
| | | https://www.bethyl.com/product/IHC-00359 Use the link above to view SDS, a current list of citations, and other product specific information. | | | |

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019

For in vitro laboratory use only. Not for any clinical, therapeutic or diagnostic use in humans or animals. Not for human or animal consumption. This product may not be resold or modified for resale without the prior written approval of Bethyl Laboratories, Inc. The information provided in this data sheet is believed to be correct but does not purport to be all-inclusive and is intended to be used as a guide. Bethyl Laboratories, Inc. shall not be liable or responsible in any way for use of either this information or the material supplied. Disposal of hazardous material may be subject to federal, state or local laws or regulations.



Detection of human NRBF2 by immunohistochemistry. *Sample:* FFPE section of human breast carcinoma. *Antibody:* Affinity purified rabbit anti-NRBF2 (Cat. No. IHC-00359) used at a dilution of 1:250. *Detection:* DAB



Detection of mouse NRBF2 by immunohistochemistry. *Sample:* FFPE section of mouse teratoma. *Antibody:* Affinity purified rabbit anti-NRBF2 (Cat. No. IHC-00359) used at a dilution of 1:250. *Detection:* DAB