

## Human HBEGF AccuSignal ELISA Kit - KOA0456

**Code:** KOA0456

**Size:** 1 Kit

**Product Description:** Human HBEGF AccuSignal ELISA Kit - KOA0456

**PhysicalState:**

<b>Label</b>	Unconjugated
<b>Gene Name</b>	HBEGF
<b>Species Reactivity</b>	Human
<b>Storage Condition</b>	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.
<b>Synonyms</b>	Diphtheria toxin receptor, diphtheria toxin receptor (heparin-binding EGF-like growth factor), diphtheria toxin receptor (heparin-binding epidermal growth factor-like growth factor), DT R, DT-R, DTR, DTS, DTSF, HB-EGF, HBEGF, HBEGF_HUMAN, HEGFL, Heparin binding EGF like growth factor, Heparin binding epidermal growth factor, Heparin binding epidermal growth factor like growth factor, Heparin-binding EGF-like growth factor, Proheparin binding EGF like growth factor
<b>Application Note</b>	Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1ml per well of the 1000pg/ml, 500pg/ml, 250pg/ml, 125pg/ml, 62.5pg/ml, 31.2pg/ml, 15.6pg/ml human HBEGF standard solutions into the precoated 96-well plate. Add 0.1ml of the sample diluent buffer into the control well (Zero well). Add 0.1ml of each properly diluted sample of human cell culture supernates, serum or plasma (heparin, EDTA) to each empty well. It is recommended that each human HBEGF standard solution and each sample be measured in duplicate.
<b>Background</b>	HBEGF (Heparin-binding EGF-like growth factor), also known as HEGFL or DTR, is a member of the EGF family of proteins that in humans is encoded by the HBEGF gene. The HBEGF gene is assigned to chromosome 5, thus confirming the assignment of the gene on the basis of its role in relation to diphtheria toxin susceptibility. HB-EGF is an 87 amino acid glycoprotein which displays highly regulated gene expression. It has been shown to play a role in wound healing, cardiac hypertrophy and heart development and function. HB-EGF binding and activation of EGF receptors plays a critical role during cardiac valve tissue development and the maintenance of normal heart function in adults.
<b>Purity And Specificity</b>	Natural and recombinant human HBEGF. There is no detectable cross-reactivity with other relevant proteins.
<b>ELISA</b>	15.6pg/ml-1000pg/ml
<b>Expiration</b>	See kit insert for complete instructions.
<b>Immunogen</b>	Expression system for standard: sf21; Immunogen sequence: D63-L148
<b>Anti-Coagulant</b>	Heparin Sodium

**Related Products**

MB-008	10X PBS pH 7.2 (0.2 M Potassium Phosphate 1.5 M Sodium Chloride) - MB-008
MB-012	10X TBS pH 7.5 (1.0 M Tris HCl 1.5 M Sodium Chloride) - MB-012
MB-013	10X TTBS pH 7.5 (1.0 M Tris HCl 1.5 M Sodium Chloride 0.1% (w/v) Tween-20) - MB-013
MB-075-1000	10X PBST (0.2 M Potassium Phosphate 1.5 M Sodium Chloride, 0.5% (v/v) Tween-20, pH 7.2) - MB-075-1000

**Related Links**

UniProtKB - Q99075

<http://www.uniprot.org/uniprot/Q99075>

 NCBI - [http://www.ncbi.nlm.nih.gov/protein/NP\\_001936.1](http://www.ncbi.nlm.nih.gov/protein/NP_001936.1)

GeneID - 1839


<http://www.ncbi.nlm.nih.gov/sites/entrez?db=gene&term=1839>

KOA0456 Protocol [http://www.rockland-inc.com/uploadedfiles/Support/KOA0456%20\(AZP\).pdf](http://www.rockland-inc.com/uploadedfiles/Support/KOA0456%20(AZP).pdf)

## Images

1

Human HBEGF AccuSignal ELISA Kit standard curve. Assay Range: 15.6pg/ml-1000pg/ml. Sensitivity: <10pg/ml.



## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.