

**Mouse CXCL5/ENA-78 AccuSignal ELISA Kit - KOA0555**
**Code:** KOA0555

**Size:** 1 Kit

**Product Description:** Mouse CXCL5/ENA-78 AccuSignal ELISA Kit - KOA0555

**PhysicalState:**

<b>Label</b>	Unconjugated
<b>Gene Name</b>	CXCL5
<b>Species Reactivity</b>	Mouse
<b>Storage Condition</b>	Store vials at 4°C prior to opening. Centrifuge product if not completely clear after standing at room temperature. This product is stable for 6 months at 4°C as an undiluted liquid. Dilute only prior to immediate use. For extended storage freeze at -20°C or below for 12 months. Avoid cycles of freezing and thawing.
<b>Synonyms</b>	AMCF II, AMCFII, C X C motif chemokine 5, Chemokine (C X C motif) ligand 5, CXCL 5, ENA 78, ENA78, Epithelial derived neutrophil activating protein 78, GCP 2, LIX, Neutrophil activating peptide ENA 78, Neutrophil activating protein 78, SCYB5, Scyb6, Small inducible cytokine B5, Small inducible cytokine subfamily B (Cys X Cys) member 5 (epithelial derived neutrophil activating peptide 78), Small inducible cytokine subfamily B, member 15
<b>Application Note</b>	Useful in Sandwich ELISA for Quantitative Detection of Antigen. Aliquot 0.1ml per well of the 1000pg/ml, 500pg/ml, 250pg/ml, 125pg/ml, 62.5pg/ml, 31.2pg/ml, 15.6pg/ml mouse CXCL5 standard solutions into the precoated 96-well plate. Add 0.1ml of the sample diluent buffer into the control well (Zero well). Add 0.1ml of each properly diluted sample of mouse cell culture supernates, serum or plasma (heparin, EDTA, citrate) to each empty well. It is recommended that each mouse CXCL5 standard solution and each sample be measured in duplicate.
<b>Background</b>	C-X-C motif chemokine 5 is a protein that in humans encoded by the CXCL5 gene. The protein encoded by this gene, CXCL5 is a small cytokine belonging to the CXC chemokine family that is also known as epithelial-derived neutrophil-activating peptide 78(ENA-78). It is produced following stimulation of cells with the inflammatory cytokines interleukin-1 or tumor necrosis factor-alpha. Expression of CXCL5 has also been observed in eosinophils, and can be inhibited with the type II interferon IFN-gamma. This chemokine stimulates the chemotaxis of neutrophils possessing angiogenic properties. It elicits these effects by interacting with the cell surface chemokine receptor CXCR2. The gene for CXCL5 is encoded on four exons and is located on human chromosome 4 amongst several other CXC chemokine genes. CXCL5 has been implicated in connective tissue remodeling. CXCL5 plays a role in reducing sensitivity to sunburn pain in some subjects, and is a potential target which can be utilized to understand more about pain in other inflammatory conditions like arthritis and cystitis.
<b>Purity And Specificity</b>	Natural and recombinant mouse CXCL5. There is no detectable cross-reactivity with other relevant proteins.
<b>ELISA</b>	15.6pg/ml-1000pg/ml
<b>Expiration</b>	See kit insert for complete instructions.
<b>Immunogen</b>	Expression system for standard: E.coli; Immunogen sequence: V45-A118
<b>Anti-Coagulant</b>	Heparin Sodium
<b>Related Products</b>	
MB-008	10X PBS pH 7.2 (0.2 M Potassium Phosphate 1.5 M Sodium Chloride) - MB-008
MB-012	10X TBS pH 7.5 (1.0 M Tris HCl 1.5 M Sodium Chloride) - MB-012
MB-013	10X TTBS pH 7.5 (1.0 M Tris HCl 1.5 M Sodium Chloride 0.1% (w/v) Tween-20) - MB-013
MB-075-1000	10X PBST (0.2 M Potassium Phosphate 1.5 M Sodium Chloride, 0.5% (v/v) Tween-20, pH 7.2) - MB-075-1000
<b>Related Links</b>	
	UniProtKB - P50228
	<a href="http://www.uniprot.org/uniprot/P50228">http://www.uniprot.org/uniprot/P50228</a>
NCBI - NP_033167.2	<a href="http://www.ncbi.nlm.nih.gov/protein/NP_033167.2">http://www.ncbi.nlm.nih.gov/protein/NP_033167.2</a>

GeneID - 20311


<http://www.ncbi.nlm.nih.gov/sites/entrez?db=gene&term=20311>

KOA0555 Protocol <http://www.rockland-inc.com/uploadedfiles/Support/KOA0555-%20SZ.pdf>

## Images

1

Mouse CXCL5/ENA-78 AccuSignal ELISA Kit standard curve.  
Assay Range: 15.6pg/ml-1000pg/ml. Sensitivity: <10pg/ml.



## Disclaimer

This product is for research use only and is not intended for therapeutic or diagnostic applications. Please contact a technical service representative for more information. All products of animal origin manufactured by Rockland Immunochemicals are derived from starting materials of North American origin. Collection was performed in United States Department of Agriculture (USDA) inspected facilities and all materials have been inspected and certified to be free of disease and suitable for exportation. All properties listed are typical characteristics and are not specifications. All suggestions and data are offered in good faith but without guarantee as conditions and methods of use of our products are beyond our control. All claims must be made within 30 days following the date of delivery. The prospective user must determine the suitability of our materials before adopting them on a commercial scale. Suggested uses of our products are not recommendations to use our products in violation of any patent or as a license under any patent of Rockland Immunochemicals, Inc. If you require a commercial license to use this material and do not have one, then return this material, unopened to: Rockland Inc., P.O. BOX 5199, Limerick, Pennsylvania, USA.