

Product Information

Product ID D3202
CAS No. 2050-87-5
Chemical Name Di-2-propenyl trisulfide

Synonym Allyl trisulfide, DATS, Allitridin

Formula C₆H₁₀S₃
Formula Wt. 178.34

Melting Point


Purity ≥98%

Solubility Soluble in acetone. Slightly soluble in ethanol (3 mg/mL), DMSO (5 mg/mL), 2-propanol, and DMF (10 mg/mL).

Store Temp -20° C

Ship Temp Ambient

Description Diallyl trisulfide is an organosulfur compound initially found in garlic; it exhibits anticancer chemotherapeutic, immunostimulatory, antioxidative, hepatoprotective, anti-fibrotic, anti-estrogenic, anti-metastatic, anti-inflammatory, anti-angiogenic, and cardioprotective activities. In vivo, diallyl trisulfide inhibits lipid peroxidation, increases phase II enzymes such as catalase, superoxide dismutase, glutathione peroxidase, and glutathione reductase, and decreases arsenic-induced hepatic dysfunction. Diallyl trisulfide also decreases collagen deposition, hepatic stellate cell activation, fibrosis, and levels of procollagen I and α-SMA. In animal models of colitis, diallyl trisulfide limits NF-κB activation, STAT3 signaling, and production of COX-2 and iNOS. Diallyl trisulfide inhibits diabetic cardiomyopathy, preventing decreases in cardiac function; it also decreases Notch1 signaling and suppresses angiogenesis in osteosarcoma cells. In breast cancer cells, diallyl trisulfide decreases expression and activity of ERB. In bladder carcinoma cells, this compound inhibits cellular migration and invasion and decreases matrix metalloproteinase (MMP) activity. Additionally, diallyl trisulfide modulates activity of HSP27. In leukemia cells, this compound induces G0/G1 phase cell cycle arrest and apoptosis and increases activation of caspase 3; in similar in vivo models, it increases macrophage phagocytosis and NK cell activity. In animal models of glioblastoma, diallyl trisulfide increases expression of p21 and p53, decreases expression of MDM2, survivin, Bcl-2, VEGF, mTOR, and c-Myc, limits activity of HDACs, and suppresses tumor growth.



Bulk quantities available upon request

Product ID	Size
D3202	100 mg
D3202	500 mg
D3202	1 g

References Suda S, Watanabe K, Tanaka Y, et al. Identification of molecular target of diallyl trisulfide in leukemic cells. *Biosci Biotechnol Biochem.* 2014;78(8):1415-7. PMID: 25130746.

Sumedha N, Miltonprabu S. Diallyl trisulfide ameliorates arsenic-induced hepatotoxicity by abrogation of oxidative stress, inflammation, and apoptosis in rats. *Hum Exp Toxicol.* 2014 Jul 25. [Epub ahead of print]. PMID: 25062976.

Hung FM, Shang HS, Tang NY, et al. Effects of diallyl trisulfide on induction of apoptotic death in murine leukemia WEHI-3 cells in vitro and alterations of the immune responses in normal and leukemic mice in vivo. *Environ Toxicol.* 2014 May 28. [Epub ahead of print]. PMID: 24890016.

Zhu X, Zhang F, Zhou L, et al. Diallyl trisulfide attenuates carbon tetrachloride-caused liver injury and fibrogenesis and reduces hepatic oxidative stress in rats. *Naunyn Schmiedebergs Arch Pharmacol.* 2014 May;387(5):445-55. PMID: 24557053.

Hahn ER, Singh SV. Diallyl trisulfide inhibits estrogen receptor-α activity in human breast cancer cells. *Breast Cancer Res Treat.* 2014 Feb;144(1):47-57. PMID: 24487688.

Shin DY, Cha HJ, Kim GY, et al. Inhibiting invasion into human bladder carcinoma 5637 cells with diallyl trisulfide by inhibiting matrix metalloproteinase activities and tightening tight junctions. *Int J Mol Sci.* 2013 Oct 1;14(10):19911-22. PMID: 24084732.

Wallace GC 4th, Haar CP, Vandergrift WA 3rd, et al. Multi-targeted DATS prevents tumor progression and promotes apoptosis in

Caution: This product is intended for laboratory and research use only. It is not for human or drug use.