

Phone: 888-558-5227

651-644-8424

Fax: 888-558-7329 Email: getinfo@lktlabs.com

Web: lktlabs.com

Product Information

Product ID N1994 CAS No. **Chemical Name**

Synonym

Formula C₇₀H₁₁₅N₂₃O₁₇

Formula Wt. 1550.84

Melting Point

Purity ≥95%

Solubility Soluble in water.

H-pGlu-Ser-His-Ile-Ser-Lys-Ala-Arg-Arg-Pro-Tyr-Ile-Leu-NH₂

Pricing and Availability

Bulk quanitites available upon request

Product ID	Size	List Price
N1994	0.5 mg	\$37.80
N1994	1 mg	\$63.00
N1994	2.5 mg	\$113.20

Store Temp -20°C Ship Temp Ambient

Description Neurotensin is a tridecapeptide that exhibits antinociceptive, pro-inflammatory, and gastrointestinal motility modulating activities. Neurotensin induces contractions in smooth muscle cells, playing a role in regulating digestive activity in vitro and in vivo. Additionally, neurotensin induces intracellular Ca2+ release, activating non-selective cation channels and altering pacemaker currents involved in gastrointestinal and colonic motility. In animal models, neurotensin displays sympathoinhibitory activity, decreases blood pressure and heart rate. Neurotensin exerts its actions through activation of neurotensin receptors; activation of these receptors also stimulates orexin neurons, suggesting a role for neurotensin in mediation of reward, feeding, and wakefulness.

References Kleczkowska P, Lipkowski AW. Neurotensin and neurotensin receptors: characteristic, structure-activity relationship and pain modulation--a review. Eur J Pharmacol. 2013 Sep 15;716(1-3):54-60. PMID: 23500196.

> Degolier TF, Brown DR, Duke GE, et al. Neurotensin and cholecystokinin contract gallbladder circular muscle in chickens. Poult Sci. 2013 Aug;92(8):2156-62. PMID: 23873564.

Furutani N, Hondo M, Kageyama H, et al. Neurotensin co-expressed in orexin-producing neurons in the lateral hypothalamus plays an important role in regulation of sleep/wakefulness states. PLoS One. 2013 Apr 19;8(4):e62391. PMID: 23620827.

Lee J, Kim YD, Park CG, et al. Neurotensin modulates pacemaker activity in interstitial cells of Cajal from the mouse small intestine. Mol Cells. 2012 May;33(5):509-16. PMID: 22441675.

Zogovic B, Pilowsky PM. Intrathecal neurotensin is hypotensive, sympathoinhibitory and enhances the baroreflex in anaesthetized rat. Br J Pharmacol. 2012 May;166(1):378-89. PMID: 22035146.

Law IK, Murphy JE, Bakirtzi K, et al. Neurotensin-induced proinflammatory signaling in human colonocytes is regulated by Barrestins and endothelin-converting enzyme-1-dependent endocytosis and resensitization of neurotensin receptor 1. J Biol Chem. 2012 Apr 27;287(18):15066-75. PMID: 22416137.

Caution: This product is intended for laboratory and research use only. It is not for human or drug use.